

Interventions for visual field defects in people with stroke

Hazelton, Christine; Campbell, Pauline; Rowe, Fiona J.; Jonuscheit, Sven; Kernohan, Ashleigh; Angilley, Jayne; Henderson, Clair A.; Langhorne, Peter; Pollock, Alex

Published in:
Stroke

DOI:
[10.1161/STROKEAHA.119.026516](https://doi.org/10.1161/STROKEAHA.119.026516)

Publication date:
2019

Document Version
Author accepted manuscript

[Link to publication in ResearchOnline](#)

Citation for published version (Harvard):

Hazelton, C, Campbell, P, Rowe, FJ, Jonuscheit, S, Kernohan, A, Angilley, J, Henderson, CA, Langhorne, P & Pollock, A 2019, 'Interventions for visual field defects in people with stroke', *Stroke*, vol. 50, no. 12, pp. e419–e420. <https://doi.org/10.1161/STROKEAHA.119.026516>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

Interventions for visual field defects in people with stroke

Keywords: stroke, visual field, quality of life, meta-analysis, literature review

Corresponding author: Christine Hazelton PhD

Nursing, Midwifery and Allied Health Professions Research Unit, A600 Govan Mbeki Building,
Glasgow Caledonian University, Cowcaddens Road, Glasgow, UK. G4 0BA

christine.hazelton@gcu.ac.uk

0141 3318181

Twitter: [@CRHazelton](https://twitter.com/CRHazelton)

Authors:

Pauline Campbell PhD, Nursing, Midwifery and Allied Health Professions Research Unit, Glasgow Caledonian University, Glasgow, UK

Fiona J Rowe PhD, Department of Health Services Research, University of Liverpool, Liverpool, UK

Sven Jonuscheit PhD, Vision Sciences, Glasgow Caledonian University, Glasgow, UK

Ashleigh Kernohan PhD, Institute of Health & Society, Newcastle University, Newcastle upon Tyne, UK

Jayne Angilley BSc, Cornwall & Isles of Scilly Primary Care Trust, Newquay, UK

Clair A Henderson MSc, Police Division, The Scottish Government, Edinburgh, UK

Peter Langhorne PhD, Academic Section of Geriatric Medicine, ICAMS, University of Glasgow, Glasgow, UK

Alex Pollock PhD, Nursing, Midwifery and Allied Health Professions Research Unit, A600 Govan Mbeki Building, Glasgow Caledonian University UK

Visual field defects (VFD) affect around one fifth of stroke survivors. They typically cause loss of up to half of an individual's vision, severely limiting the sensory information they receive about their environment. VFDs impact on practical aspects of daily life, social activities are negatively affected, and psychological effects include loss of self-confidence. A range of interventions exist: these are proposed to work through restitution of the lost visual field; compensation via behaviour changes; substitution using a device or extraneous modification; or assessment/screening, ensuring diagnosis and treatment.

In this Cochrane review we aimed to determine the effects of any intervention targeting visual field defects in people with stroke.

Methods We searched 9 databases including MEDLINE, Embase, CINAHL, and clinical trials databases to May 2018. We also searched reference lists, hand-searched journals and conference proceedings, and contacted experts. We included randomised trials in adults after stroke (and mixed populations): the primary outcome was ability in activities of daily living (ADL). Two reviewers independently screened abstracts, extracted data and appraised risk of bias due to allocation concealment, masking, incomplete outcome data and other sources. Outcome data were pooled in meta-analyses, and quality assessed using the GRADE approach.

Main results We included 20 studies (n=547/732; stroke/total participants), with a mean of 36 stroke participants (range 10-87). Eleven studies compared an intervention with placebo, control, no treatment or standard care: three explored restitution, four compensation, three substitution and one assessment/screening.

Restitution: very low quality evidence showed that visual restitution therapy had a beneficial effect on quality of life (one study, n=19, Odds Ratio (OR) 13.00, 95% CI 2.07 to 81.48), but no effect on visual field. Limitations with these data however mean it is not possible to draw a clear conclusion about therapy effectiveness.

Compensation: low quality evidence showed scanning training improved quality of life (two studies, n=96, mean difference (MD) 9.36, 95% CI 3.10 to 15.62) (Fig 1). There was low or very-low quality evidence of no effect on visual field, extended ADL, reading, and scanning ability. Low-quality evidence also showed no significant increase in adverse events.

Substitution: very low-quality evidence suggested no effect on ADL, extended ADL, quality of life or reading. Prisms benefited scanning ability (one study, n=39, MD 9.80, 95% CI 1.91 to 17.69) but with increased odds of an adverse event (primarily headache)(one study, n=59, OR 87.32, 95% CI 4.87 to 1564.66).

Assessment/Screening: very low-quality evidence suggested that assessment by an orthoptist had no effect on ADL.

Conclusions We found limited evidence relating to VFD in stroke: the effectiveness of interventions is still not clear. Whilst our analysis suggests scanning training may benefit quality of life, further research is likely to alter this result.

Implications for practice Scanning training may be effective in improving quality of life for stroke survivors with VFD, without risk of side effects. However, it is not clear if other approaches are effective, and the use of prisms may lead to minor adverse events.

Implications for research Adequately powered randomised controlled studies that compare an intervention with appropriate control are needed to determine effectiveness.

Words 509

Figure 1. Forest plot of compensatory interventions (scanning training) versus control, placebo or no intervention, for quality of life outcomes

(attached as a separate file)

Disclosures: Fiona Rowe was the chief investigator for the VISION trial (Rowe 2010), funded by the Stroke Association: Alex Pollock was also involved in this study. Fiona Rowe was a co-investigator for the Jarvis 2012 trial, funded by the University of Liverpool. Christine Hazelton has carried out non-randomised studies in to the effectiveness of a number of scanning training interventions, including the intervention studied by Roth 2009, and is developing further project proposals in this area. No other interests are known.

Funding: Alex Pollock and Pauline Campbell are employed by the Nursing, Midwifery and Allied Health Professions Research Unit, which is funded by the Chief Scientist Office in Scotland. Royal National Institute of Blind People (RNIB, Scotland) funded the first (2009) version of this review.

Acknowledgement: This paper is based on a Cochrane Review published in The Cochrane Library 2019, Issue 5 (see www.thecochranelibrary.com for information). Cochrane Reviews are regularly updated as new evidence emerges and in response to feedback, and The Cochrane Library should be consulted for the most recent version of the review.

Pollock A, Hazelton C, Rowe FJ, Jonuscheit S, Kernohan A, Angilley J, Henderson CA, Langhorne P, Campbell P. Interventions for visual field defects in people with stroke. Cochrane Database of Systematic Reviews 2019, Issue 5. Art. No.: CD008388. DOI: 10.1002/14651858.CD008388.pub3.

<http://dx.doi.org/10.1002/14651858.CD008388>